

Chapter 7: Moving Beyond Linearity^e

So far we have mainly focused on linear models.

Linear models are simple to describe and implement.

Advantage: interpretation/inference.

disadvantage: can have limited predictive performance because linearity is always an approximation.

Previously, we have seen we can improve upon least squares using ridge regression, the lasso, principal components regression, and more.

improvement obtained by reducing complexity of OLS \Rightarrow lowering variance.

Still a linear model! Can only be improved so much.

Through simple and more sophisticated extensions of the linear model, we can relax the linearity assumption while still maintaining as much interpretability as possible. \rightarrow extensions of linear model.

① Polynomial regression: adding extra predictors that are original variables raised to a power.

e.g. cubic regression uses X, X^2, X^3 as predictors, e.g. $y = \beta_0 + \beta_1 X + \beta_2 X^2 + \beta_3 X^3 + \epsilon$.

+ non-linear fit

- large powers can lead to strange shapes (especially near the boundary).

② Step functions: cut the range of a variable into K distinct regions to produce a categorical variable. Fit a piecewise constant function to X .

③ Regression Splines: more flexible than polynomials + step functions (extends both).

idea: cut the range of X into K distinct regions + fit polynomial within each region

Polynomials are constrained so that they are smoothly joined.

④ Generalized additive models (GAM): extends above to deal w/ multiple predictors.

We will start w/ predicting Y on X ($p=1$) and extend to multiple.

Note: We can talk regression or classification, e.g. logistic regression = $P(Y=1|X) = \frac{\exp(\beta_0 + \beta_1 X + \beta_2 X^2 + \dots + \beta_d X^d)}{1 + \exp(\beta_0 + \beta_1 X + \dots + \beta_d X^d)}$

We've seen this one.

1 Step Functions

Using polynomial functions of the features as predictors imposes a global structure on the non-linear function of X .

We can instead use *step-functions* to avoid imposing a global structure.

i.e. break range of X into bins and fit a different constant in each bin.

details: ① create cut points C_1, \dots, C_k in the range of X .

② Construct $K+1$ new variables

$$C_0(X) = \mathbb{I}(X < C_1)$$

$$C_1(X) = \mathbb{I}(C_1 \leq X < C_2)$$

\vdots

$$C_K(X) = \mathbb{I}(C_K \leq X)$$

} indicator functions
"dummy variables"

Note for any X ,

$$C_0(X) + C_1(X) + \dots + C_K(X) = 1.$$

since X must be in exactly 1 interval.

← leave out $C_0(X)$ because it is equivalent to fitting an intercept.

③ Use OLS to fit linear model using $C_1(X), \dots, C_K(X)$

$$Y = \beta_0 + \beta_1 C_1(X) + \dots + \beta_K C_K(X) + \varepsilon.$$

For a given value of X , at most one of C_1, \dots, C_K can be non-zero.

When $X < C_1 \Rightarrow$ all predictors $C_1, \dots, C_K = 0$

$\Rightarrow \beta_0$ interpreted as mean value for Y when $X < C_1$.

β_j represents the average increase in mean response for $X \in [C_j, C_{j+1})$ relative to $X < C_1$.

We can also fit the logistic regression model for classification:


$$P(Y=1|X) = \frac{\exp(\beta_0 + \beta_1 C_1(X) + \dots + \beta_K C_K(X))}{1 + \exp(\beta_0 + \beta_1 C_1(X) + \dots + \beta_K C_K(X))}$$

← new interpretation of β_j 's related to log-odds.

Example: Wage data.

$n=3000$ male workers in Mid-atlantic region.

year	age	maritl	race	education	region	jobclass	health	health_ins	logwage	wage
2006	18	1. Never Married	1. White	1. < HS Grad	2. Middle Atlantic	1. Industrial	1. <=Good	2. No	4.318063	75.04315
2004	24	1. Never Married	1. White	4. College Grad	2. Middle Atlantic	2. Information	2. >=Very Good	2. No	4.255273	70.47602
2003	45	2. Married	1. White	3. Some College	2. Middle Atlantic	1. Industrial	1. <=Good	1. Yes	4.875061	130.98218
2003	43	2. Married	3. Asian	4. College Grad	2. Middle Atlantic	2. Information	2. >=Very Good	1. Yes	5.041393	154.68529



fitted value of wage using step function of age.

missed increasing trend here.

logistic regression modeling prob. of being a high earner (Wage > 250k).

using step function w/ knots at $x=30, 60$.

Unless there are natural breakpoints in the predictor, piecewise constant functions can miss trends.

2 Basis Functions

Polynomial and piecewise-constant regression models are in fact special cases of a *basis function approach*.

Idea:

have a family of functions or transformations that can be applied to a variable X

$$b_1(x), b_2(x), \dots, b_K(x).$$

Instead of fitting the linear model in X , we fit the model

$$y_i = \beta_0 + \beta_1 b_1(x_i) + \dots + \beta_K b_K(x_i) + \varepsilon_i$$

Note that the basis functions are fixed and known. (we choose them).

e.g. Polynomial regression: $b_j(x_i) = x_i^j$, $j = 1, \dots, d$

e.g. Step functions: $b_j(x_i) = \mathbb{I}(c_j \leq x_i < c_{j+1})$.

We can think of this model as a standard linear model with predictors defined by the basis functions and use least squares to estimate the unknown regression coefficients.

\Rightarrow We can use all our inference tools for linear models, e.g. $se(\hat{\beta}_j)$ and F -statistics for model significance.

Many alternatives exist for basis functions:

e.g. wavelets, Fourier series, regression splines (next).

3 Regression Splines

Regression splines are a very common choice for basis function because they are quite flexible, but still interpretable. Regression splines extend upon polynomial regression and piecewise constant approaches seen previously.

start with

3.1 Piecewise Polynomials

Instead of fitting a high degree polynomial over the entire range of X , piecewise polynomial regression involves fitting separate low-degree polynomials over different regions of X .

i.e. fit two different polynomials to data: 1 on subset for $x < c$ and a second on subset for $x \geq c$.
↓ single cut-point.

For example, a piecewise cubic with no knots is just a standard cubic polynomial.

A piecewise cubic with a single knot at point c takes the form

$$y_i = \begin{cases} \beta_{01} + \beta_{11}x_i + \beta_{21}x_i^2 + \beta_{31}x_i^3 + \varepsilon_i & \text{if } x_i < c \\ \beta_{02} + \beta_{12}x_i + \beta_{22}x_i^2 + \beta_{32}x_i^3 + \varepsilon_i & \text{if } x_i \geq c \end{cases}$$

each polynomial can be fit using least squares.

Using more knots leads to a more flexible piecewise polynomial.

If we place L knots \Rightarrow fit $L+1$ polynomials (doesn't have to be cubic).

In general, we place L knots throughout the range of X and fit $L + 1$ polynomial regression models.

This leads to $(d+1)(L+1)$ degrees of freedom in the model
(# parameters to fit \approx complexity/flexibility).

3.2 Constraints and Splines

To avoid having **too much flexibility**, we can **constrain** the piecewise polynomial so that the fitted curve must be continuous.

i.e. there cannot be a jump at the knots.

To go further, we could add two more constraints


- ① first derivatives of the piecewise polynomials are continuous at the knots
- ② 2nd derivatives of the piecewise polynomials are continuous at the knots.

In other words, we are requiring the piecewise polynomials to be **smooth**.


Each constraint that we impose on the piecewise cubic polynomials effectively frees up one degree of freedom, by reducing the complexity of the resulting fit.

The fit with continuity and smoothness constraints is called a spline.


A degree- d spline is a piecewise degree- d polynomial w/ continuity in derivatives up to degree $d-1$ at each knot.



piecewise cubic polynomial.



piecewise cubic with continuity



*cubic spline
cts + cts 1st and 2nd derivatives (smooth).*

3.3 Spline Basis Representation

Fitting the spline regression model is more complex than the piecewise polynomial regression. We need to fit a degree d piecewise polynomial and also constrain it and its $d - 1$ derivatives to be continuous at the knots.

up to

We can use the basis model to represent a regression spline.

e.g. cubic
Spline w/ L knots:

$$y_i = \beta_0 + \beta_1 b_1(x_i) + \beta_2 b_2(x_i) + \dots + \beta_{L+3} b_{L+3}(x_i) + \varepsilon_i$$

for appropriate functions b_1, \dots, b_{L+3} .

x, x^2, x^3

The most direct way to represent a cubic spline is to start with the basis for a cubic polynomial and add one truncated power basis function per knot.

$$h(x, c) = (x - c)_+^3 = \begin{cases} (x - c)^3 & \text{if } x > c \\ 0 & \text{o.w.} \end{cases} \quad \text{where } c \text{ is the knot.}$$

$$\Rightarrow y_i = \beta_0 + \beta_1 x_i + \beta_2 x_i^2 + \beta_3 x_i^3 + \sum_{j=1}^L \beta_{3+j} h(x_i, c_j) + \varepsilon_i$$

see homework

→ This will lead to discontinuity in only the 3rd derivative at each c_j with continuous first and second derivatives and continuity at each c_j .

df : $L + 4$ (cubic spline w/ L knots).

Unfortunately, splines can have high variance at the outer range of the predictors. One solution is to add boundary constraints.

i.e. X is small or large.

require function to be linear at the boundary (where x is smaller than the smallest knot or bigger than the biggest knot)

"natural spline"

additional constraint produces more stable estimates at the boundaries.

3.4 Choosing the Knots

When we fit a spline, where should we place the knots?

Regression spline is most flexible in regions that contain a lot of knots (coefficients can change more rapidly).
 \Rightarrow place knots where we think relationship changes rapidly (less stable).

More common in practice: place them uniformly

to do this, choose desired degrees of freedom (flexibility) + use software to automatically place knots at uniform quantiles of the data.

How many knots should we use?

\Leftrightarrow how many degrees of freedom should we use?

Use CV! Choose L that gives smallest CV error!


alternative: penalized splines (splines + lasso).

3.5 Comparison to Polynomial Regression

Regression splines often gives superior results to polynomial regression.

Polynomial regression must use high degree to achieve flexible fit (e.g. x^{15}),

but regression splines introduce flexibility through knots (but fixed degree) \Rightarrow more stability (esp. at boundaries).



extra flexibility of polynomial produces undesirable result at the boundary but spline w/ same flexibility still reasonable.

4 Generalized Additive Models

So far we have talked about flexible ways to predict Y based on a single predictor X .

These approaches can be seen as extensions of simple linear regression

$$Y = \beta_0 + \beta_1 X + \varepsilon.$$

Generalized Additive Models (GAMs) provide a general framework for extending a standard linear regression model by allowing non-linear functions of each of the variables while maintaining **additivity**.

flexibly predicting Y on the basis of several predictors X_1, \dots, X_p

4.1 GAMs for Regression — still additive models

↳ can also be used for classification using logistic regression.

A natural way to extend the multiple linear regression model to allow for non-linear relationships between feature and response:

linear regression: $y_i = \beta_0 + \beta_1 x_{i1} + \dots + \beta_p x_{ip} + \varepsilon_i$

idea: replace each linear component $\beta_j x_{ij}$ with a smooth non-linear function:

$$\Rightarrow \text{GAM: } g_i = \beta_0 + \sum_{j=1}^p f_j(x_{ij}) + \varepsilon_i$$

$$= \beta_0 + f_1(x_{i1}) + f_2(x_{i2}) + \dots + f_p(x_{ip}) + \varepsilon_i$$

"additive" because we calculate a separate f_j for each X_j and add them together.

possibilities for f_j :

- identity (leads to linear regression).
- polynomial
- regression splines
- smoothing splines, local linear regression. → see textbook ch. 7.5–7.6

The beauty of GAMs is that we can use our fitting ideas in this chapter as building blocks for fitting an additive model.

Example: Consider the Wage data.


$$\text{Wage} = \beta_0 + f_1(\text{year}) + f_2(\text{age}) + f_3(\text{education}) + \varepsilon$$

where f_1 is natural spline w/ 4 df

f_2 is natural spline w/ 5 df

f_3 is identity of dummy variables created from education.

easy to fit least squares by choosing appropriate basis functions.



Pros and Cons of GAMs

4.2 GAMs for Classification

GAMs can also be used in situations where Y is categorical. Recall the logistic regression model:

A natural way to extend this model is for non-linear relationships to be used.

Example: Consider the Wage data.

